

ON THE DEFINITION OF THE KOBAYASHI-BUSEMAN PSEUDOMETRIC

NIKOLAI NIKOLOV AND PETER PFLUG

ABSTRACT. We prove that the $(2n-1)$ -th Kobayashi pseudometric of any domain $D \subset \mathbb{C}^n$ coincides with the Kobayashi–Buseman pseudometric of D , and that $2n-1$ is the optimal number, in general.

1. INTRODUCTION AND RESULTS

Let $\mathbb{D} \subset \mathbb{C}$ be the unit disc. Recall first the definitions of the Lempert function $\tilde{k}_D$ and the Kobayashi–Royden pseudometric k_D of a domain $D \subset \mathbb{C}^n$ (cf. [1]):

$$\begin{aligned}\tilde{k}_D(z, w) &= \inf\{\tanh^{-1}|\alpha| : \exists \varphi \in \mathcal{O}(\mathbb{D}, D) : \varphi(0) = z, \varphi(\alpha) = w\}, \\ \kappa_D(z; X) &= \inf\{\alpha \geq 0 : \exists \varphi \in \mathcal{O}(\mathbb{D}, D) : \varphi(0) = z, \alpha\varphi'(0) = X\},\end{aligned}$$

where $z, w \in D$, $X \in \mathbb{C}^n$. The Kobayashi pseudodistance k_D can be defined as the largest pseudodistance which does not exceed $\tilde{k}_D$. Note that if $k_D^{(m)}$ denotes the m -th Lempert function of D , that is,

$$k_D^{(m)}(z, w) = \inf\left\{\sum_{j=1}^m \tilde{k}_D(z_j, z_{j+1}) : z_1, \dots, z_m \in D, z_1 = z, z_m = w\right\},$$

then

$$k_D(z, w) = \inf_m k_D^{(m)}(z, w) = \inf\left\{\int_0^1 \kappa_D(\gamma(t); \gamma'(t)) dt\right\},$$

where the infimum is taken over all piecewise C^1 -curves $\gamma : [0, 1] \rightarrow D$ connecting z and w . By a result of M. Y. Pang (see [5]), the Kobayashi–Royden pseudometric is the infinitesimal form of the Lempert function

2000 *Mathematics Subject Classification.* 32F45.

Key words and phrases. Lempert function, Kobayashi pseudodistance, Kobayashi–Royden pseudometric, Kobayashi–Buseman pseudometric.

This note was written during the stay of the first named author at the Universität Oldenburg supported by a grant from the DFG (January – March 2006). He likes to thank both institutions for their support.

for taut domains; more precisely, if D is a taut domain, then

$$(1) \quad \kappa_D(z; X) = \lim_{\mathbb{C}_* \ni t \rightarrow 0} \frac{\tilde{k}_D(z, z + tX)}{t}.$$

In [3], S. Kobayashi introduces a new invariant pseudometric, called the Kobayashi–Buseman pseudometric in [1]. One of the equivalent ways to define the Kobayashi–Buseman pseudometric $\hat{\kappa}_D$ of D is just to set $\hat{\kappa}_D(z; \cdot)$ to be largest pseudonorm which does not exceed $\kappa_D(z; \cdot)$. Recall that

$$\hat{\kappa}_D(z; X) = \inf \left\{ \sum_{j=1}^m \kappa_D(z; X_j) : m \in \mathbb{N}, \sum_{j=1}^m X_j = X \right\}.$$

Thus, it is natural to consider the new function $\kappa_D^{(m)}$, namely,

$$\kappa_D^{(m)}(z; X) = \inf \left\{ \sum_{j=1}^m \kappa_D(z; X_j) : \sum_{j=1}^m X_j = X \right\}.$$

We call $\kappa_D^{(m)}$ the m -th Kobayashi pseudometric of D . It is clear that $\kappa_D^{(m)} \geq \kappa_D^{(m+1)}$ and if $\kappa_D^{(m)}(z; \cdot) = \kappa_D^{(m+1)}(z; \cdot)$ for some m , then $\kappa_D^{(m)}(z; \cdot) = \kappa_D^{(j)}(z; \cdot)$ for any $j > m$. It is shown in [3] that

$$(2) \quad \kappa_D^{(2n)} = \hat{\kappa}_D.$$

Let now $D \subset \mathbb{C}^n$ be a taut domain. We point out that, using the equalities (1) and (2), M. Kobayashi (see [2]) shows that

$$\hat{\kappa}_D(z; X) = \lim_{\mathbb{C}_* \ni t \rightarrow 0} \frac{k_D(z, z + tX)}{t}.$$

Obvious modifications in the proof of this result lead to

$$(3) \quad \lim_{u, v \rightarrow z, u \neq v} \frac{k_D^{(m)}(u, v) - \kappa_D^{(m)}(z; u - v)}{\|u - v\|} = 0.$$

uniformly in m and locally uniformly in z ; thus,

$$\kappa_D^{(m)}(z; X) = \lim_{\mathbb{C}_* \ni t \rightarrow 0} \frac{k_D^{(m)}(z, z + tX)}{t}$$

uniformly in m and locally uniformly in z and X .

The aim of this note is the following result which improves (2).

Theorem 1. *For any domain $D \subset \mathbb{C}^n$ we have that*

$$(4) \quad \kappa_D^{(2n-1)} = \hat{\kappa}_D.$$

On the other hand, if $n \geq 2$ and

$$D_n = \{z \in \mathbb{C}^n : \sum_{j=2}^n (2|z_1^3 - z_j^3| + |z_1^3 + z_j^3|) < 2(n-1)\},$$

then

$$(5) \quad \kappa_{D_n}^{(2n-2)}(0; \cdot) \neq \hat{\kappa}_{D_n}(0; \cdot).$$

Note that the proof below shows that the equality (4) remains true for any n -dimensional complex manifold.

An immediate consequence of Theorem 1 and the equality (3) is:

Corollary 2. *For any taut domain $D \subset \mathbb{C}^n$ one has that*

$$\lim_{w \rightarrow z, w \neq z} \frac{k_D^{(2n-1)}(z, w)}{k_D(z, w)} = 1$$

locally uniformly in z , and $2n-1$ is the optimal number, in general.

Remarks. (i) If $D \subset \mathbb{C}$, then even $\tilde{k}_D = k_D$ (cf. [1]).

(ii) Corollary 2 holds for n -dimensional taut complex manifolds.

(iii) Observe that Corollary 2 may be taken as a very weak version of the following question asked by S. Krantz (see [4]): whether there is a positive integer $m = m(D)$ such that $k_D = k_D^{(m)}$.

Let now h_S be the Minkowski functions of a starlike domain $S \subset \mathbb{R}^N$, that is, $h_S(X) = \inf\{t > 0 : X/t \in S\}$. We may define as above

$$h_S^{(m)}(X) = \inf\left\{\sum_{j=1}^m h_S(X_j) : \sum_{j=1}^m X_j = X\right\}.$$

Then the Minkowski function $h_{\hat{S}}$ of the convex hull $\hat{S}$ of S is the largest pseudonorm which does not exceed h_S . It follows by a lemma due to C. Carathéodory (cf. [2]) that

$$(6) \quad h_{\hat{S}} = h_S^{(N)} = \inf\left\{\sum_{j=1}^M h_S(X_j) : M \leq N, \sum_{j=1}^M X_j = X, \right. \\ \left. X_1, \dots, X_M \text{ are } \mathbb{R}\text{-linearly independent}\right\}.$$

One can easily see that N is the optimal number for the class of starlike domains in $\mathbb{R}^N$.

Denote by $I_{D,z}$ the indicatrix of $\kappa_D(z; \cdot)$, that is, $I_{D,z} = \{X \in \mathbb{C}^n : \kappa_D(z; X) < 1\}$. Note that $I_{D,z}$ is a balanced domain (a domain $B \subset \mathbb{C}^n$ is said to be balanced if $\lambda X \in B$ for any $\lambda \in \overline{\mathbb{D}}$ and any $X \in B$). In particular, $I_{D,z}$ is a starlike domain and hence (2) follows by (6). Similarly, (4) will follow by the following.

Proposition 3. *If $B \subset \mathbb{C}^n$ is a balanced domain, then*

$$(7) \quad h_{\hat{B}} = h_B^{(2n-1)}.$$

Observe that the domain D_n from Theorem 1 is pseudoconvex and balanced, thus $\kappa_{D_n}(0; \cdot) = h_{D_n}$ (cf. [1]) and so $\kappa_{D_n}^{(m)}(0; \cdot) = h_{D_n}^{(m)}$. Then inequality (5) is equivalent to

$$(8) \quad h_{\hat{D}_n} \neq h_{D_n}^{(2n-2)}.$$

2. PROOFS

To prove Proposition 3, we shall need the following result.

Lemma 4. *Any balanced domain can be exhausted by bounded balanced domains with continuous Minkowski functions.*

Proof. Let $B \subset \mathbb{C}^n$ be a balanced domain. Denote by $\mathbb{B}_n(z, r) \subset \mathbb{C}^n$ the ball with center z and radius r . For $z \in \mathbb{C}^n$ and $j \in \mathbb{N}$, set $F_{n,j,z} := \overline{\mathbb{B}_n(z, \|z\|^2/j)}$. We may assume that $\mathbb{B}_n(0, 1) \subset \subset B$. Put

$$B_j := \{z \in \mathbb{B}_n(0, j) : F_{n,j,z} \subset B\}, \quad j \in \mathbb{N}.$$

Then $(B_j)_{j \in \mathbb{N}}$ is an exhaustion of B by non-empty bounded open sets. We shall show that B_j is a balanced domain with continuous Minkowski functions.

For this, take any $z \in B_j$ and $0 \neq \lambda \in \overline{\mathbb{D}}$, and observe that $F_{n,j,\lambda z} \subset \lambda F_{n,j,z} \subset B$. Thus, B_j is a balanced domain.

Since h_{B_j} is an upper semicontinuous function, it remains to prove that it is lower semicontinuous. Assuming the contrary, we may find a sequence of points $(z_k)_{k \in \mathbb{N}}$ converging to some point $z \in \mathbb{C}^n$ and a positive number c such that $h_{B_j}(z_k) < 1/c < h_{B_j}(z)$ for any k . Note that $F_{n,j,cz_k} \subset B$, $k \in \mathbb{N}$. Hence $\mathbb{B}_n(cz, c^2\|z\|^2/j) \subset B$. On the other hand, fix $t \in (0, 1)$ such that $h_{B_j}(tcz) > 1$. Then $F_{n,j,tcz} \subset \mathbb{B}_n(cz, c^2\|z\|^2/j) \subset B$; thus $h_{B_j}(tcz) < 1$, a contradiction. $\square$

Proof of Proposition 3. First, we shall prove (7) in the case, when $B \subset \mathbb{C}^n$ is a bounded balanced domain with continuous Minkowski function. Fix a vector $X \in \mathbb{C}^n \setminus \{0\}$. Then $h_{\hat{B}}(X) \neq 0$ and we may assume that $h_{\hat{B}}(X) = 1$. By the continuity of h_B and (6), there exist $\mathbb{R}$ -linearly independent vectors $X_1, \dots, X_m$ ($m \leq 2n$) such that $\sum_{j=1}^m X_j = X$ and

$\sum_{j=1}^m h_B(X_j) = 1$. Since $h_{\hat{B}}$ is a norm, the triangle inequality implies that $h_B(X_j) = h_{\hat{B}}(X_j)$, $j = 1, \dots, m$. To prove (7), it suffices to show

that $m \neq 2n$. The convexity of $\hat{B}$ provides a support hyperplane H for $\hat{B}$ at $X \in \partial\hat{B}$, say $H = \{z \in \mathbb{C}^n : \operatorname{Re}\langle z - X, \overline{X_0} \rangle = 0\}$, $X_0 \in \mathbb{C}^n$, where $\langle \cdot, \cdot \rangle$ stands for the Hermitian scalar product in $\mathbb{C}^n$. Assuming

$$m = 2n \text{ implies that } H = \left\{ \sum_{j=1}^m \alpha_j \hat{X}_j : \sum_{j=1}^m \alpha_j = 1, \alpha_1, \dots, \alpha_m \in \mathbb{R} \right\},$$

where $\hat{X}_j := X_j/h_B(X_j) \in \partial\hat{B}$. In particular, $\partial\hat{B}$ contains a relatively open subset of H . Since $\hat{B}$ is a balanced domain, it follows that its intersection with the plane, spanned by X_0 , is a disc whose boundary contains a line segment, a contradiction.

Now let $B \subset \mathbb{C}^n$ be an arbitrary balanced domain. If $(B_j)_{j=1}^\infty$ is an exhaustion of B given by Lemma 4, then $h_{B_j} \searrow h_B$ pointwise and hence $h_{\hat{B}_j} \searrow h_{\hat{B}}$ by (6). Then (7) follows by the inequalities $h_{\hat{B}} \leq h_B^{(2n-1)} \leq h_{B_j}^{(2n-1)}$ and the equality $h_{\hat{B}_j} = h_{B_j}^{(2n-1)}$ from above. $\square$

Proof of the inequality (8). Let $L_n = \{z \in \mathbb{C}^n : z_1 = 1\}$. Then the triangle inequality implies that $D_n \subset \mathbb{D} \times \mathbb{C}^{n-1}$ and

$$F_n := \partial D_n \cap L_n = \{z \in \mathbb{C}^n : z_1 = 1, z_j \in \Omega, 2 \leq j \leq n\},$$

where Ω is the set of the third roots of unity. Denoting by Δ the convex hull of Ω , it follows that

$$\partial\hat{D}_n \cap L_n = \hat{F}_n = \{1\} \times \Delta^{n-1}.$$

Hence, $\partial\hat{D}_n \cap L_n$ is a $(2n-2)$ -dimensional convex set. Put $\tilde{F}_n = \{Y \in \hat{F}_n : h_{D_n}^{(2n-2)}(Y) = 1\}$. If $X \in \tilde{F}_n$, then there exist $X_1, \dots, X_m \in \mathbb{C}^n \setminus \{0\}$, $m \leq 2n-2$, such that $\sum_{j=1}^m X_j = X$ and $\sum_{j=1}^m h_{D_n}(X_j) = 1$ (note that D_n is taut). Hence, $X_1/h_{D_n}(X_1), \dots, X_m/h_{D_n}(X_m) \in F_n$ and X belongs to their convex hull. Since F_n is a finite set, it follows that $\tilde{F}_n$ is contained in a finite union of at most $(2n-3)$ -dimensional convex sets. Thus, $\hat{F}_n \neq \tilde{F}_n$ which implies that $h_{\hat{D}_n} \neq h_{D_n}^{(2n-2)}$. $\square$

REFERENCES

- [1] M. Jarnicki, P. Pflug, *Invariant distances and metrics in complex analysis*, de Gruyter Exp. Math. 9, de Gruyter, Berlin, New York, 1993.
- [2] M. Kobayashi, *On the convexity of the Kobayashi metric on a taut complex manifold*, Pacific J. Math. 194 (2000), 117–128.
- [3] S. Kobayashi, *A new invariant infinitesimal metric*, Internat. J. Math. 1 (1990), 83–90.
- [4] S. Krantz, *Convexity in complex analysis*, Several Complex Variables and Complex Geometry, Proc. Symp. Pure Math., vol. 52, Amer. Math. Soc., Providence, RI, 1991, pp. 119–137.

- [5] M.-Y. Pang, *On infinitesimal behavior of the Kobayashi distance*, Pacific J. Math. 162 (1994), 121–141.

INSTITUTE OF MATHEMATICS AND INFORMATICS, BULGARIAN ACADEMY OF SCIENCES, ACAD. G. BONCHEV 8, 1113 SOFIA, BULGARIA

E-mail address: nik@math.bas.bg

CARL VON OSSIEZKY UNIVERSITÄT OLDENBURG, FACHBEREICH MATHEMATIK, POSTFACH 2503, D-26111 OLDENBURG, GERMANY

E-mail address: pflug@mathematik.uni-oldenburg.de